



APPLN. FILING DATE: AUGUST 22, 2001  
TITLE: METHOD FOR MAKING MORPHOLINO-NUCLEOTIDES,  
AND THEIR USE FOR ANALYSING AND MARKING NUCLEIC  
ACID SEQUENCES  
INVENTOR(S): MARCIACQ, FLORENCE ET. AL.  
APPLN. SERIAL NO.: 09/914,221

## Figure 5

Sequence to be analysed (polymerase template)

3' HO-ApTpGpCpApApTpCpCpGpApTpGpApCpTpGpApGpCpCpApTpCpG 5'  
(SEQ ID NO: 1)

Primer


+

5' TpApCpGpTpTpApGpGpC-OH 3' (SEQ ID NO: 2)



3' HO-ApTpGpCpApApTpCpCpGpApTpGpApCpTpGpApGpCpCpApTpCpG 5'  
5' TpApCpGpTpTpApGpGpC-OH 3'

DNA Polymerase



5' Thymidine triphosphate






Figure 6

Sequence to be analysed (polymerase template)

3' HO-ApTpGpCpApApTpCpCpGpApTpGpApCpTpGpApGpCpCpApTpCpG 5'  
(SEQ ID NO: 1)

Primer


+

5' TpApCpGpTpTpApGpGpC-OH 3' (SEQ ID NO: 2)




3' HO-ApTpGpCpApApTpCpCpGpApTpGpApCpTpGpApGpCpCpApTpCpG 5'  
5' TpApCpGpTpTpApGpGpC-OH 3'

DNA Polymerase



3'-deoxythymidine 5'-triphosphate



The newly synthesized chain no longer contains 3' OH  
required to attach the next nucleotide